

RESEARCH ARTICLE

Risk factors during first 1,000 days of life for carotid intima-media thickness in infants, children, and adolescents: A systematic review with meta-analyses

Adina Mihaela Epure^{1,2,*}, Magali Rios-Leyvraz², Daniela Anker³, Stefano Di Bernardo⁴, Bruno R. da Costa^{3,5}, Arnaud Chiolerio^{1,2,3,6}, Nicole Sekarski⁴

1 Population Health Laboratory (#PopHealthLab), University of Fribourg, Fribourg, Switzerland, **2** Department of Epidemiology and Health Services, Center for Primary Care and Public Health (UNISANTÉ), University of Lausanne, Lausanne, Switzerland, **3** Institute of Primary Health Care (BIHAM), University of Bern, Bern, Switzerland, **4** Paediatric Cardiology Unit, Woman-Mother-Child Department, Lausanne University Hospital (CHUV), Lausanne, Switzerland, **5** Institute of Health Policy, Management and Evaluation, University of Toronto, Toronto, Canada, **6** Department of Epidemiology, Biostatistics and Occupational Health, McGill University, Montréal, Canada

✉ These authors contributed equally to this work.

* adina-mihaela.epure@unifr.ch



OPEN ACCESS

Citation: Epure AM, Rios-Leyvraz M, Anker D, Di Bernardo S, da Costa BR, Chiolerio A, et al. (2020) Risk factors during first 1,000 days of life for carotid intima-media thickness in infants, children, and adolescents: A systematic review with meta-analyses. PLoS Med 17(11): e1003414. <https://doi.org/10.1371/journal.pmed.1003414>

Academic Editor: Sanjay Basu, Harvard Medical School, UNITED STATES

Received: May 6, 2020

Accepted: October 19, 2020

Published: November 23, 2020

Peer Review History: PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pmed.1003414>

Copyright: © 2020 Epure et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data and citations to all data sources, that is, the full-text

Abstract

Background

The first 1,000 days of life, i.e., from conception to age 2 years, could be a critical period for cardiovascular health. Increased carotid intima-media thickness (CIMT) is a surrogate marker of atherosclerosis. We performed a systematic review with meta-analyses to assess (1) the relationship between exposures or interventions in the first 1,000 days of life and CIMT in infants, children, and adolescents; and (2) the CIMT measurement methods.

Methods and findings

Systematic searches of Medical Literature Analysis and Retrieval System Online (MEDLINE), Excerpta Medica database (EMBASE), and Cochrane Central Register of Controlled Trials (CENTRAL) were performed from inception to March 2019. Observational and interventional studies evaluating factors at the individual, familial, or environmental levels, for instance, size at birth, gestational age, breastfeeding, mode of conception, gestational diabetes, or smoking, were included. Quality was evaluated based on study methodological validity (adjusted Newcastle–Ottawa Scale if observational; Cochrane collaboration risk of bias tool if interventional) and CIMT measurement reliability. Estimates from bivariate or partial associations that were least adjusted for sex were used for pooling data across studies, when appropriate, using random-effects meta-analyses. The research protocol was published and registered on the International Prospective Register of Systematic Reviews (PROSPERO; CRD42017075169). Of 6,221 reports screened, 50 full-text articles from 36 studies (34 observational, 2 interventional) totaling 7,977 participants (0 to 18 years at CIMT assessment) were retained. Children born small for gestational age had increased CIMT (16

publications of included studies, are within the manuscript and its [supporting information](#) files.

Funding: This work was funded by the Swiss National Science Foundation (www.snf.ch; project number 32003B-163240; grantee: AC). The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Abbreviations: ART, assisted reproductive technologies; CCA, common carotid artery; CENTRAL, Cochrane Central Register of Controlled Trials; CI, confidence interval; CIMT, carotid-intima media thickness; CVD, cardiovascular disease; DOHaD, Developmental Origins of Health and Disease; EMBASE, Excerpta Medica database; MEDLINE, Medical Literature Analysis and Retrieval System Online; MOOSE, Meta-analysis of Observational Studies in Epidemiology; PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-Analyses; PROSPERO, International Prospective Register of Systematic Reviews; SMD, standardized mean difference.

studies, 2,570 participants, pooled standardized mean difference (SMD): 0.40 (95% confidence interval (CI): 0.15 to 0.64, p : 0.001), I^2 : 83%). When restricted to studies of higher quality of CIMT measurement, this relationship was stronger (3 studies, 461 participants, pooled SMD: 0.64 (95% CI: 0.09 to 1.19, p : 0.024), I^2 : 86%). Only 1 study evaluating small size for gestational age was rated as high quality for all methodological domains. Children conceived through assisted reproductive technologies (ART) (3 studies, 323 participants, pooled SMD: 0.78 (95% CI: -0.20 to 1.75, p : 0.120), I^2 : 94%) or exposed to maternal smoking during pregnancy (3 studies, 909 participants, pooled SMD: 0.12 (95% CI: -0.06 to 0.30, p : 0.205), I^2 : 0%) had increased CIMT, but the imprecision around the estimates was high. None of the studies evaluating these 2 factors was rated as high quality for all methodological domains. Two studies evaluating the effect of nutritional interventions starting at birth did not show an effect on CIMT. Only 12 (33%) studies were at higher quality across all domains of CIMT reliability. The degree of confidence in results is limited by the low number of high-quality studies, the relatively small sample sizes, and the high between-study heterogeneity.

Conclusions

In our meta-analyses, we found several risk factors in the first 1,000 days of life that may be associated with increased CIMT during childhood. Small size for gestational age had the most consistent relationship with increased CIMT. The associations with conception through ART or with smoking during pregnancy were not statistically significant, with a high imprecision around the estimates. Due to the large uncertainty in effect sizes and the limited quality of CIMT measurements, further high-quality studies are needed to justify intervention for primordial prevention of cardiovascular disease (CVD).

Author summary

Why was this study done?

- Exposure to adverse experiences in the first 1,000 days of life, i.e., from conception to age 2 years, may determine adaptative changes in the blood vessel walls and increased carotid intima-media thickness (CIMT) in infants, children, and adolescents. This implies carotid arteries with thicker walls that may be due to changes in blood flow and pressure or other factors related to the process of atherosclerosis.

What did the researchers do and find?

- We performed a systematic review of published studies with meta-analyses and included 36 studies, involving 7,977 participants between 0 and 18 years at CIMT assessment.
- Risk factors in the first 1,000 days of life, particularly poor fetal growth, are associated with increased CIMT in infants, children, and adolescents.
- There is scarce evidence from interventional studies beginning in the first 1,000 days, and none was shown to prevent or improve vascular remodeling in children.
- CIMT measurement protocols in children are heterogeneous and often poorly reported.

What do these findings mean?

- From a public health perspective, acting early in life by preventing risk factors such as poor fetal growth could help maintain a low cardiovascular risk over the life course.
- Assessing vascular structure and function in children is important to better characterize lifetime risk trajectories and tailor primordial prevention of cardiovascular disease (CVD). Primordial prevention aims to prevent the development of risk factors instead of treating them.
- From a clinical standpoint, promotion of a healthy lifestyle is important at any age, and screening of postnatal cardiovascular risk factors, targeted at children exposed to risk factors in the first 1,000 days of life, may be warranted.
- A widely accepted standardized CIMT measurement protocol in children is needed.

Introduction

Background

Within a Developmental Origins of Health and Disease (DOHaD) framework, the first 1,000 days of life, i.e., the period from conception to age 2 years, is regarded as a critical period for long-term cardiovascular health [1]. Evidence suggests that adaptations in body structure and function during development, as a response to cues from the environment, for instance, assisted reproductive technologies (ART) for conception [2], undernutrition, or environmental pollutants [1], could increase the risk for cardiovascular disease (CVD). Indeed, several studies showed that a low birth weight is associated with higher blood pressure [3] or diabetes risk [4], as well as with cardiovascular mortality in adulthood [5].

Carotid intima-media thickness (CIMT) is a marker of CVD risk linked to pathways that originate in the first 1,000 days of life. In adults, CIMT is predictive of heart attack and stroke risk [6,7] and is also associated with prenatal risk factors, such as impaired fetal growth and preterm birth [8,9]. Within a DOHaD perspective, these associations may be explained by early life adaptative changes in the vascular phenotype [10], with lifelong effects on the CVD risk. Nonetheless, in children, the evidence on increased CIMT after impaired fetal growth or preterm birth is inconsistent [11–14]. Further, CVD risk is determined by a combination of factors, at multiple levels, rather than a single factor [15]. A large range of early life factors at the child, familial, or environmental levels needs to be explored to better understand CIMT changes in children. To our knowledge, such a comprehensive overview is lacking.

CIMT assessment in children can be challenging [16,17]. Several aspects related to the ultrasound equipment [18–20], the site of measurement, or the edge detection approach may influence the reliability of measurements [17,21]. Further, measurements in infants and young children are challenging due to limited compliance, anatomic particularities, and a lack of recommendations tailored to this age group [16,22]. The comparability of studies is therefore limited, and the inconsistency in findings may be attributed, at least in part, to the heterogeneity in CIMT measurement. An appraisal of the CIMT measurement methods in children is needed to shed light on these limitations.

We therefore aimed to perform a systematic review with meta-analyses to (1) assess the relationship between exposures or interventions in the first 1,000 days of life and CIMT in infants, children, and adolescents; and (2) critically appraise the CIMT measurement methods.

Methods

Protocol development and reporting

The protocol for this systematic review was registered on the International Prospective Register of Systematic Reviews (PROSPERO; CRD42017075169) and published in full in a peer-reviewed journal [23]. The reporting of this paper complies with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines [24] ([S1 PRISMA Checklist](#)) and the Meta-analysis of Observational Studies in Epidemiology (MOOSE) guidelines [25].

Eligibility criteria

Study designs. Observational and interventional studies with the following designs were considered for inclusion: cohort, case-control, cross-sectional studies, randomized trials, non-randomized controlled trials, and noncontrolled trials. Case reports, case series, opinion papers, letters to the editor, comments, conference abstracts, policy papers, reviews and meta-analyses, study protocols without baseline data, and animal studies were excluded.

Participants. We considered for inclusion studies in children from birth up to 18 years old, including studies where participants were recruited pre-birth. Apparently, healthy children and participants with clinical conditions relating to the first 1,000 days of life (e.g., prematurity) were included. Studies in children with rare or special conditions, such as congenital heart diseases, or postnatal conditions conferring an increased risk of CVD, such as hypertension or diabetes, were not considered for inclusion. We aimed to make inferences to the general population. Further, we assumed that there may be a different effect of DOHAD exposures in these groups than in the general population, which should be evaluated separately.

Exposures/Interventions. We considered for inclusion prenatal and postnatal exposures, at the individual, familial, or environmental level, which reflected the developmental milieu of the child and occurred between conception and age 24 months (i.e., first 1,000 days of life). These included biological factors related to physiology, physiopathology, or epigenetics and contextual factors related to socioeconomic status, behaviors, or environmental toxins. The selection of exposures was informed by the review of Hanson and Gluckman [1] and Moore and colleagues [26] and was conditioned by the timing of occurrence or ascertainment of exposure, as appropriate. Exposures that started in the first 1,000 days of life, but continued in later life, were included if they spanned up to 5 years of age. Likewise, periconceptional factors that reflected the environment in which the baby was conceived and grew were included. For interventional studies, we considered both pharmacological and non-pharmacological interventions, at the individual, familial, or environmental level, conducted in the first 1,000 days of life. Specifically, eligible interventions had to start and end in the first 1,000 days of life or start in the first 1,000 days of life and continue in later life no longer than child age 5 years.

Comparators. Where applicable, the use of a reference group without the exposure or intervention of interest was sufficient for inclusion.

Outcome measures. The outcome was the intima-media thickness of the carotid artery measured by ultrasonography from birth to 18 years of age.

Time frame and setting. There was no restriction by length of follow-up. There was no restriction by type of setting.

Language. Studies in English and French were considered for inclusion.

Search strategy

Systematic searches were conducted in the Medical Literature Analysis and Retrieval System Online (MEDLINE) database, Excerpta Medica database (EMBASE), and Cochrane Central Register of Controlled Trials (CENTRAL) from inception to 18 March 2019. The strategies for the systematic searches included the 2 main concepts of this systematic review: (1) children and adolescents; and (2) CIMT (detailed search strategies are available in [S1 Table](#)). Supplementary searches consisted of (1) a manual search of reference lists and other reviews on the topic; (2) forward citation tracking on Web of Science based on retrieved eligible reports; and (3) personalized search queries in Google Scholar and trial registers. The full strategies are available in [S2 Table](#).

Study selection process

Following systematic searches in the databases, all retrieved reports were imported into Endnote (version X8.1, Clarivate Analytics, London, United Kingdom), and duplicates were removed according to the method of Bramer and colleagues [27]. Subsequently, reports were uploaded to Covidence (www.covidence.org), a systematic review management platform. Two reviewers performed study eligibility screening independently and in duplicate. Each reviewer screened the reports in a 2-step approach: (1) based on titles and abstracts; and (2) based on full-text of reports retained in the first step. The reviewers agreed on 94% of the titles and abstracts and on 84% of the full texts. Disagreements between reviewers were resolved by discussion, or, if necessary, by a third reviewer. The investigator of 1 completed study (ClinicalTrials.gov identifier: NCT02147457), with unpublished data on CIMT, was contacted by e-mail, but data could not be made available due to technical and logistical issues when performing measurements.

Data extraction

Data were extracted independently and in duplicate by 2 reviewers using a standardized electronic form in Microsoft Excel (version 2016, Microsoft Corporation, Redmond, United States of America). The data extraction process was piloted on a subset of articles, and a guide for extraction was designed to minimize errors and enhance consistency between reviewers. We extracted information about (1) study characteristics; (2) population characteristics; (3) CIMT definition and additional characteristics of the CIMT measurement methods; (4) characteristics of each exposure or intervention type; and (5) adjusted and unadjusted association or effect estimates.

The quality of each study was evaluated independently by 2 reviewers using an adjusted version of the Newcastle–Ottawa Scale for observational studies [28] and Cochrane collaboration risk of bias tool for interventional studies [29,30] ([S7 Table](#)). The quality of the CIMT measurement method was examined according to a predefined tool with 3 levels (higher, lower, and unclear reliability). The complete tool and the algorithm of judgment are available in [S3 Table](#).

Disagreements in extracted data between the 2 reviewers were resolved by discussion or with the arbitration of a third reviewer. Essential missing information was searched by checking additional references related to that study, such as the published research protocol. We also attempted to contact authors of 2 studies by e-mail.

Data analysis

Analyses were conducted as prespecified in the study protocol [23]. Data transformations and estimations were done according to recommendations and formulae provided in Borenstein

and colleagues [31,32], the Cochrane Handbook for Systematic Reviews [29], Lipsey and colleagues [33], Rücker and colleagues [34], Wan and colleagues [35], Aloe and colleagues [36], and Nieminen and colleagues [37]. The following data transformations were performed: (1) when the mean was not available, it was estimated from the median [35]; (2) when standard deviations were not available, they were estimated from the range or interquartile range [36], standard errors [33], and confidence intervals (CIs) [29,33]; and (3) when standard errors were not provided, they were calculated from CIs, *p*-values, and *t*-values [29,33]. To perform meta-analyses, additional data simplifications were performed as follows: (1) if a study reported on the same exposure or outcome type at multiple time points, the latest time point was included in the analyses; (2) if a study reported on multiple exposed or reference subgroups, they were combined into 1 using formulae provided by the Cochrane collaboration in case of means and standard deviations [29] or fixed-effect meta-analysis in case of association estimates for each subgroup [32]; (3) if a study provided association estimates separately for left and right CIMT or multiple sample groups, they were combined using fixed-effect meta-analysis [32,34]; (4) if a study reported on both mean and maximum CIMT, mean CIMT was included in the analyses; and (5) if multiple publications from the same study reported on the same exposure type, the publication providing data on the longest follow-up, left and right and/or mean thickness CIMT, or with the clearest reporting or highest sample size was used.

Standardized association measures were used to pool results across studies so that the comparison of different exposure types would be possible on the same scale. The standardized mean difference (SMD) in CIMT between exposed and reference groups was used as the common association measure for binary exposure variables [31–33]. When means and standard deviations for exposed and reference were not available or could not be estimated from other statistics as reported above, SMD were estimated from the unstandardized regression coefficient and the full-sample standard deviation [33,38]. The correlation coefficient was used as the common association measure for continuous exposure variables [32,33,36,37]. When the correlation was not available, it was estimated from the regression coefficient and its standard error or the unstandardized regression coefficient and the standard deviations of the exposure and outcome variables [36,37]. If the reported association measures could not be transformed to SMD or correlation coefficient, they were analyzed and reported separately. Estimates from bivariate associations or partial associations that were least adjusted for sex were used for pooling data across studies [39]. If at least 3 studies reported on the same exposure type, with a similar definition, association estimates were pooled using the DerSimonian–Laird random-effects model. As recommended by Amrhein and colleagues [40], together with more than 800 cosignatories of the initiative “Retire statistical significance,” and because we assessed multiple associations, we interpreted the point estimate as the most compatible value of the observed association. The uncertainty around the point estimate is comprised in the accompanying 95% CI, which is reported along the corresponding *p*-value.

The heterogeneity was assessed using the Cochran Q test and I^2 and τ^2 statistics [29,32,41]. I^2 and τ^2 are standard measures of heterogeneity in random-effect meta-analyses that reflect the amount of true heterogeneity between studies, i.e., the variation in effect sizes that is not due to sampling error. I^2 is the percentage of variation in effect sizes attributable to heterogeneity, ranging from 0% to 100%, with 0% indicating that all variation in effect sizes is due to sampling error within studies and 100% indicating that all variation is due to true heterogeneity between studies [32]. τ^2 is used to assign study weights under the random-effects model [32]. Sources of heterogeneity were explored using subgroup meta-analyses, meta-regression, and sensitivity analyses. Publication bias was evaluated by the visual inspection of funnel plots and, where appropriate, by Egger test. All statistical analyses were conducted with Stata (version 16, StataCorp, Texas, USA) and Microsoft Excel (version 2019).

Results

Study characteristics

The searches retrieved 8,338 reports (Fig 1). Following deduplication, titles and abstracts of 6,221 reports were screened, and 154 were selected for full-text screening. A total of 50 full-text articles, pertaining to 36 studies, were eventually included. Details on study characteristics and associated reports are presented in Table 1. Studies were conducted in Europe ($n = 28$), Asia ($n = 3$), North America ($n = 3$), and Australia ($n = 2$). Some 23 studies were conducted in healthcare facilities, 6 were community/population based, 4 included a mix of the aforementioned 2 settings, and 3 did not report any information about their participant recruitment sites or sources. Exposures were evaluated in studies with a cohort ($n = 22$) or cross-sectional ($n = 14$) design. Interventions were studied in 2 randomized controlled trials. All studies




Fig 1. Flow diagram showing the study selection process. CENTRAL, Cochrane Central Register of Controlled Trials; EMBASE, Excerpta Medica database; MEDLINE, Medical Literature Analysis and Retrieval System Online.

<https://doi.org/10.1371/journal.pmed.1003414.g001>

Table 1. Characteristics of included studies.

Author, publication year	Country	Design	Setting	N* participants	Age at CIMT assessment	Mean (SD), CIMT, mm	Exposure/intervention categories at child (1), family (2), or environmental (3) levels
Observational studies							
Gale and colleagues [42], 2006	UK	cohort	healthcare facility	216	9 years	right: 0.34 (0.06)	Diet and feeding practices (1), Fetal growth (1), Postnatal growth (1), Maternal weight and growth, nutrition, and physical activity (2), Socioeconomic status (3), Tobacco exposure (3)
Gale and colleagues [43], 2008	UK	cohort	healthcare facility	178	9 years	right: 0.34 (0.06)	Maternal weight and growth, nutrition, and physical activity (2)
Ayer and colleagues [44], 2009a	Australia	cohort	healthcare facility	405	8 years	left and right: 0.59 (0.07)	Fetal growth (1), Pregnancy-specific factors (2), Tobacco exposure (3)
Ayer and colleagues [45], 2009b	Australia	cohort	healthcare facility	405	8 years	left and right: 0.59 (0.06)	Diet and feeding practices (1)
Ayer and colleagues [46], 2011	Australia	cohort	healthcare facility	405	8 years	left and right: 0.59 (0.1)	Tobacco exposure (3)
Skilton and colleagues [47], 2012	Australia	cohort	healthcare facility	363	8 years	left and right: 0.59 (0.06)	Fetal growth (1), Gestational age (1)
Skilton and colleagues [48], 2013	Australia	cohort	healthcare facility	395	8 years	left or right: 0.77 (0.08)	Diet and feeding practices (1), Fetal growth (1), Gestational age (1), Postnatal growth (1), Socioeconomic status (3)
Crispi and colleagues [11], 2010	Spain	cohort	healthcare facility	200	2 to 6 years	left and right: 0.38 (0.11)	Fetal growth (1), Gestational age (1)
Rodriguez-Lopez and colleagues [49], 2016	Spain	cohort	healthcare facility	202	4 to 5 years	left and right: 0.38 (0.03)	Diet and feeding practices (1), Fetal growth (1), Gestational age, Other (1), Postnatal growth (1), Pregnancy-specific factors (2), Socioeconomic status (3), Tobacco exposure (3)
Trevisanuto and colleagues [50], 2010	Italy	cohort	unclear	38	3 to 5 years	left: 0.51 (0.08); right: 0.51 (0.07)	Fetal growth (1)
Evelein and colleagues [51], 2011	the Netherlands	cohort	community/population based	296	5 years	right: 0.39 (0.04)	Diet and feeding practices (1)
Geerts and colleagues [52], 2012	the Netherlands	cohort	community/population based	258	5 years	right: 0.38 (0.03)	Tobacco exposure (3)
Evelein and colleagues [53], 2013	the Netherlands	cohort	community/population based	323	5 years	right: 0.39 (0.69)	Fetal growth (1), Postnatal growth (1)
Pluymen and colleagues [54], 2017	the Netherlands	cohort	community/population based	413	5 years	right: 0.39 (0.04)	Diet and feeding practices (1)
Atabek and colleagues [55], 2011	Turkey	cohort	healthcare facility	55	48 to 72 hours	right: 0.31 (0.03)	Cardio-metabolic and inflammatory factors (1), Fetal growth (1), Pregnancy-specific factors (2)
Dratva and colleagues [12], 2013	USA	cohort	community/population based	670	mean age: 11 years	right: 0.57 (0.04)	Fetal growth (1), Gestational age (1), Pregnancy-specific factors (2)
Breton and colleagues [56], 2016b	USA	cohort	community/population based	240	mean age: 11 years	N/S: 0.55 (0.05)	Epigenetics (1)

(Continued)

Table 1. (Continued)

Author, publication year	Country	Design	Setting	N* participants	Age at CIMT assessment	Mean (SD), CIMT, mm	Exposure/intervention categories at child (1), family (2), or environmental (3) levels
Breton and colleagues [57], 2016a	USA	cohort	community/population based	392	mean age: 11 years	left: 0.56 (0.05); right: 0.57 (0.04)	Epigenetics (1), Air pollution (3)
Schubert and colleagues [58], 2013	Sweden	cohort	healthcare facility	50	3 months	N/S: 0.39 (0.12)	Gestational age (1)
Valenzuela-Alcaraz and colleagues [59], 2019	Spain	cohort	healthcare facility	160	3 years	left and right: 0.49 (0.08)	Pregnancy-specific factors (2)
Lee and colleagues [60], 2014	Germany	cohort	healthcare facility	47	10 to 14 years	left and right: 0.45 (0.03)	Gestational age (1)
Morsing and colleagues [61], 2014	Sweden	cohort	healthcare facility	93	6 to 9 years	N/S: 0.34 (0.06)	Fetal growth (1), Gestational age (1)
Stergiotou and colleagues [62], 2014	Spain	cohort	healthcare facility	201	0 to 1 week	left and right: 0.24 (0.05)	Cardio-metabolic and inflammatory factors (1), Fetal growth (1)
Gruszfeld and colleagues [63], 2015	Belgium, Germany, Italy, Poland, and Spain	cohort	unclear	383	5 years	left and right: 0.4 (0.11)	Diet and feeding practices (1), Fetal growth (1), Maternal weight and growth, nutrition, and physical activity (2), Paternal factors (2), Tobacco exposure (3)
Sebastiani and colleagues [64], 2016	Spain	cohort	healthcare facility	46	6 years	left and right: 0.35 (0.08)	Fetal growth (1)
Liu and colleagues [65], 2017	Australia	cohort	community/population based	1,477	11 to 12 years	right: 0.58 (0.05)	Socioeconomic status (3)
Mohlkert and colleagues [66], 2017	Sweden	cohort	community/population based	235	6 years	left: 0.38 (0.04); right: 0.38 (0.04)	Gestational age (1)
Tzschoppe and colleagues [67], 2017	Germany	cohort	healthcare facility	20	6 years	left and right: 0.38 (0.08)	Fetal growth (1)
Carreras-Badosa and colleagues [68], 2018	Spain	cohort	healthcare facility	66	5 to 6 years	right: 0.37 (0.03)	Maternal weight and growth, nutrition, and physical activity (2)
Chen and colleagues [69], 2019	Canada	cohort	healthcare facility	119	1 year	left and right: 0.56 (0.06)	Cardio-metabolic and inflammatory factors (1), Cardio-metabolic and inflammatory factors (2)
Prins-Van Ginkel and colleagues [70], 2019	the Netherlands	cohort	other [†]	221	16 years	left and right: 0.47 (0.04)	Cardio-metabolic and inflammatory factors (1)
Sebastiani and colleagues [71], 2019	Spain	cohort	healthcare facility	68	2 years	left and right: 0.3 (0.02)	Fetal growth (1)
Sundholm and colleagues [72], 2019	Finland	cohort	other [†]	201	mean age: 6 years	left and right: 0.36 (0.04)	Maternal weight and growth, nutrition, and physical activity (2), Pregnancy-specific factors (2)
Jouret and colleagues [13], 2011	France	cross-sectional	healthcare facility	113	mean age: 11 years	right: 0.46 (0.05)	Fetal growth (1)

(Continued)

Table 1. (Continued)

Author, publication year	Country	Design	Setting	N* participants	Age at CIMT assessment	Mean (SD), CIMT, mm	Exposure/intervention categories at child (1), family (2), or environmental (3) levels
Scherrer and colleagues [73], 2012	Switzerland	cross-sectional	other [†]	39	mean age: 11 years	left and right: 0.39 (0.03)	Pregnancy-specific factors (2)
de Arriba and colleagues [74], 2013	Spain [‡]	cross-sectional	unclear	269	4 to 15 years	N/S: 0.36 (0.08)	Fetal growth (1)
Maurice and colleagues [75], 2014	Canada	cross-sectional	healthcare facility	14	13 to 15 years	right: 0.48 (0.05)	Fetal growth (1)
Xu and colleagues [76], 2014	China	cross-sectional	healthcare facility	124	3 to 7 years	left: 0.39 (0.05)	Pregnancy-specific factors (2)
Putra and colleagues [77], 2015	Indonesia	cross-sectional	community/population based	285	15 to 18 years	N/S: 0.44 (0.07)	Diet and feeding practices (1)
Sodhi and colleagues [78], 2015	India	cross-sectional	healthcare facility	100	0 to 4 days	left and right: 0.42 (0.05)	Cardio-metabolic and inflammatory factors (1), Fetal growth (1)
Ciccone and colleagues [79], 2016	Italy	cross-sectional	healthcare facility	30	3 to 5 years	left: 0.51 (0.06); right: 0.52 (0.06)	Fetal growth (1), Gestational age (1)
Faienza and colleagues [80], 2016	Italy	cross-sectional	healthcare facility	52	mean age: 10 years	left and right: 0.48 (0.06)	Fetal growth (1)
Olander and colleagues [81], 2016	Finland	cross-sectional	healthcare facility	174	5 to 184 hours	left and right: 0.17 (0.03)	Cardio-metabolic and inflammatory factors (1), Fetal growth (1), Gestational age (1)
Dilli and colleagues [82], 2017	Turkey	cross-sectional	healthcare facility	60	0 to 48 hours	left and right: 0.32 (0.03)	Cardio-metabolic and inflammatory factors (1), Fetal growth (1)
Stock and colleagues [14], 2018	Austria and Italy	cross-sectional	community/population based	930	15 to 16 years	left and right: 0.38 (0.05)	Fetal growth (1), Gestational age (1)
Wilde and colleagues [83], 2018	the Netherlands	cross-sectional	other [†]	162	6 to 8 years	right: 0.38 (0.04)	Other (2)
Muñiz Fontán and colleagues [84], 2019	Spain	cross-sectional	healthcare facility	239	6 to 8 years	left and right: 0.43 (0.04)	Fetal growth (1)
Interventional studies							
Ayer and colleagues [45], 2009b	Australia	randomized controlled trial	healthcare facility	405	8 years	left and right: 0.59 (0.06)	Diet and feeding practices (1)
Gruszfeld and colleagues [63], 2015	Belgium, Germany, Italy, Poland, and Spain	randomized controlled trial	unclear	383	5 years	left and right: 0.4 (0.11)	Diet and feeding practices (1)

Note: Studies reporting on both interventions and exposures were analyzed as randomized controlled trial for the data on interventions and as observational cohort for the data on exposures ($N = 2$, namely Ayer and colleagues [45], 2009b and Gruszfeld and colleagues [63], 2015).

*N = number of participants with CIMT measurements introduced in the analyses.

[†]Mix of healthcare facility and community based.

[‡]Country of the first author.

CIMT, carotid intima-media thickness; N/S, not specified; SD, standard deviation; UK, United Kingdom; USA, United States of America.

<https://doi.org/10.1371/journal.pmed.1003414.t001>

comprised girls and boys. Overall, the minimum age at CIMT assessment was 5 hours, and the maximum age was 18 years.

CIMT measurement methods

Image acquisition methods were relatively uniform across studies, but image analysis methods varied greatly (Tables 2 and S4). Measurements were performed on the common carotid artery (CCA) in the majority of studies (92%). However, poor or missing information on key characteristics of the CIMT measurement, namely the carotid wall, the method for edge detection and analysis of the distance between interfaces, or quality control procedures were relatively common. The segmental thickness (mean or maximum) and the type and number of sides (right, left, or the mean of both) included in the CIMT calculation were among the main sources of heterogeneity between studies. Several studies used multiple approaches for analysis; thus, up to 3 different CIMT outcomes were reported in each publication. Only 12 (33.3%) studies were at higher quality across all domains of CIMT reliability.

Risk factors at the child level

Child risk factors evaluated across studies included (1) small size for gestational age and other fetal growth indicators, such as birth weight, length, head circumference; (2) prematurity; or (3) duration of breastfeeding. The effect of interventions targeting the dietary intake of proteins and polyunsaturated fatty acids after birth was also assessed.

CIMT was higher in children born small for gestational age (16 studies, 2,570 participants, pooled SMD 0.40 (95% CI: 0.15 to 0.64, p : 0.001), I^2 : 83%) compared with those with an appropriate size at birth (Table 3; Fig 2A). The same relationship was found when analyses were restricted to children born small and with a documented prenatal diagnosis of intrauterine growth restriction (IUGR; 8 studies, 623 participants, pooled SMD 0.35 (95% CI: -0.06 to 0.77, p : 0.097), I^2 : 77%). There was no clear association of prematurity with CIMT when data were pooled across all studies (7 studies, 2,024 participants, pooled SMD 0.03 (95% CI: -0.17 to 0.22, p : 0.805), I^2 : 50%) (Table 3; Fig 3A). However, small size for gestational age (3 studies, 461 participants, pooled SMD: 0.64 (95% CI: 0.09 to 1.19, p : 0.024), I^2 : 86%) (Fig 2B) and prematurity (3 studies, 736 participants, pooled SMD: 0.25 (95% CI: 0.02 to 0.48, p : 0.032), I^2 : 0%) (Fig 3B) were consistently associated with higher CIMT, and the associations increased in magnitude when restricted to studies of higher CIMT quality. Additionally, subgroup analyses showed that CIMT was higher in infants than in older children exposed to these risk factors (Table 4). Meta-analyses of other fetal growth indicators showed that CIMT tended to be lower with higher values of birth weight or length, but the magnitude of these relationships was low (Table 3).

Inconclusive evidence was found for breastfeeding. Four studies identified a lower CIMT among children with a higher duration of breastfeeding (S5 Table). Two studies identified a higher CIMT among children exclusively breastfed compared to exclusively formula fed (S5 Table). Meta-analysis was not possible due to very heterogeneous comparisons across studies. Two interventional studies on dietary interventions at the child level involved a total of 787 participants. In 1 study, a diet comprising a higher n-3 to n-6 fatty acids ratio from birth to age 5 years had no effect on CIMT (SMD 0 (95% CI: -0.20 to 0.20)) [45]. In the other study, non-breastfed children on formulas with higher protein content during their first year of life had a lower CIMT (SMD -0.21 (95% CI: -0.45 to 0.04)) compared to children on lower protein formulas [63]. Scarce evidence, reported in 1 study, existed for other factors, such as cardio-metabolic and inflammatory factors, for instance, blood pressure [81], cord blood cholesterol [55], cortisol [69], and epigenetic changes, such as DNA methylation levels following exposure to air pollution [56,57] (Tables 1 and S6).

Table 2. CIMT measurement characteristics.

Author, publication year	N outcomes	Image acquisition			Image analysis			Reliability		
		Side	Segment	Wall	Edge detection, analysis of the distance between interfaces	Cardiac cycle phase	Wall thickness	Acquisition site	Analysis	Reproducibility assessment
Gale and colleagues [41,43], 2006, 2008	1	right	CCA	far	N/S, automatic over a specific length	end diastole	N/S	higher	higher	higher
Ayer and colleagues [44–46], 2009a, 2009b, 2011; Skilton and colleagues [47], 2012	1	left and right	CCA	far	automatic/ semiautomatic, automatic over a specific length	end diastole	mean	higher	higher	higher
Skilton and colleagues [85], 2013	1	left or right	CCA	far	automatic/ semiautomatic, automatic over a specific length	end diastole	maximum	higher	higher	higher
Crispi and colleagues [11,85], 2010, 2012	1	left and right	CCA	far	automatic/ semiautomatic, automatic over a specific length	end diastole	mean	higher	higher	higher
Rodriguez-Lopez and colleagues [49], 2016	1	left and right	CCA	far	automatic/ semiautomatic, automatic over a specific length	end diastole	N/S	higher	higher	higher
Trevisanuto and colleagues [50], 2010	2	left; right	CCA	far	manual, point-to-point measurements	N/S	maximum	higher	lower	higher
Evelein and colleagues [51,53], 2011, 2013 Geerts and colleagues [52], 2012	1	right	CCA	far	automatic/ semiautomatic, automatic over a specific length	end diastole	N/S	higher	higher	higher
Pluymen and colleagues [54], 2017	1	right	CCA	far	automatic/ semiautomatic, automatic over a specific length	end diastole	maximum	higher	higher	higher
Atabek and colleagues [55], 2011	1	right	CCA	far	manual, point-to-point measurements	N/S	maximum	higher	lower	unclear
Dratva and colleagues [12], 2013	1	right	CCA	far	automatic/ semiautomatic, automatic over a specific length	end diastole	mean	higher	higher	higher
Breton and colleagues [57], 2016a	2	left; right	CCA	far	automatic/ semiautomatic, automatic over a specific length	end diastole	N/S	higher	higher	higher
Breton and colleagues [56], 2016b	1	N/S	CCA	far	automatic/ semiautomatic, automatic over a specific length	end diastole	N/S	higher	higher	higher
Schubert and colleagues [58], 2013	1	N/S	CCA	far	N/S, automatic over a specific length	end diastole	mean	higher	higher	higher
Valenzuela-Alcaraz and colleagues [59,86], 2013, 2019	2	left and right	CCA	far	automatic/ semiautomatic, automatic over a specific length	end diastole	mean; maximum	higher	higher	higher
Lee and colleagues [60], 2014	1	left and right	CCA	far	automatic/ semiautomatic, automatic over a specific length	N/S	mean	higher	higher	higher

(Continued)

Table 2. (Continued)

Author, publication year	N outcomes	Image acquisition			Image analysis			Reliability		
		Side	Segment	Wall	Edge detection, analysis of the distance between interfaces	Cardiac cycle phase	Wall thickness	Acquisition site	Analysis	Reproducibility assessment
Morsing and colleagues [61], 2014	1	N/S	CCA	N/S	manual, point-to-point measurements	end diastole	mean	unclear	lower	unclear
Stergiotou and colleagues [62], 2014	2	left and right	CCA	far	automatic/ semiautomatic, automatic over a specific length	end diastole	mean; maximum	higher	higher	higher
Gruszfeld and colleagues [63], 2015	1	left and right	CCA	far	manual, point-to-point measurements	end diastole	mean	higher	lower	lower
Sebastiani and colleagues [64], 2016	1	left and right	CCA	far	N/S, N/S	end diastole	N/S	higher	unclear	higher
Liu and colleagues [65], 2017	1	right	CCA	far	automatic/ semiautomatic, automatic over a specific length	end diastole	maximum	higher	higher	higher
Mohlkert and colleagues [66], 2017	2	left; right	CCA	far	automatic/ semiautomatic, automatic over a specific length	end diastole	mean	higher	higher	unclear
Tzschoppe and colleagues [67], 2017	1	left and right	CCA	N/S	N/S, N/S	N/S	maximum	unclear	unclear	unclear
Carreras-Badosa and colleagues [68], 2018	1	right	CCA	N/S	N/S, N/S	N/S	N/S	unclear	unclear	higher
Chen and colleagues [69], 2019	2	left and right; left or right	other*	far	N/S, automatic over a specific length	N/S	N/S; maximum	unclear	unclear	unclear
Prins-Van Ginkel and colleagues [70], 2019	1	left and right	CCA	N/S	automatic/ semiautomatic, automatic over a specific length	end diastole	N/S	unclear	higher	unclear
Sebastiani and colleagues [71], 2019	1	left and right	CCA	far	N/S, N/S	end diastole	N/S	higher	unclear	higher
Sundholm and colleagues [72], 2019	1	left and right	CCA	far	manual, N/S	end diastole	N/S	higher	unclear	higher
Jouret and colleagues [13], 2011	1	right	CCA	N/S	N/S, automatic over a specific length	N/S	N/S	unclear	unclear	unclear
Scherrer and colleagues [73,87], 2012	1	left and right	CCA	N/S	automatic/ semiautomatic, automatic over a specific length	end diastole	mean	unclear	higher	higher
de Arriba and colleagues [74], 2013	1	N/S	CCA	far	N/S, N/S	N/S	maximum	higher	unclear	unclear
Maurice and colleagues [75], 2014	1	right	CCA	far	automatic/ semiautomatic, automatic over a specific length	end diastole	N/S	higher	higher	unclear
Xu and colleagues [76], 2014	1	left	CCA	N/S	N/S, N/S	N/S	N/S	unclear	unclear	unclear
Putra and colleagues [77], 2015	1	N/S	N/S	N/S	N/S, N/S	N/S	N/S	unclear	unclear	unclear
Sodhi and colleagues [78], 2015	2	left and right	CCA	N/S	manual, point-to-point measurements	N/S	mean; maximum	unclear	lower	lower

(Continued)

Table 2. (Continued)

Author, publication year	N outcomes	Image acquisition			Image analysis			Reliability		
		Side	Segment	Wall	Edge detection, analysis of the distance between interfaces	Cardiac cycle phase	Wall thickness	Acquisition site	Analysis	Reproducibility assessment
Ciccone and colleagues [79], 2016	2	left; right	CCA	far	N/S, N/S	end diastole	N/S	higher	unclear	unclear
Faienza and colleagues [80], 2016	1	left and right	CCA	N/S	manual, point-to-point measurements	end diastole	N/S	unclear	lower	higher
Olander and colleagues [81], 2016	1	left and right	CCA	far	manual, point-to-point measurements	end diastole	mean	higher	lower	higher
Dilli and colleagues [82], 2017	3	left; right; left and right	CCA	far	N/S, automatic over a specific length	end diastole	N/S	higher	higher	higher
Stock and colleagues [14], 2018	2	left and right; left or right	CCA	far	manual, point-to-point measurements	N/S	mean; maximum	higher	lower	unclear
Wilde and colleagues [83], 2018	1	right	CCA	far	automatic/semiautomatic, automatic over a specific length	diastole	N/S	higher	higher	higher
Muñiz Fontán and colleagues [84], 2019	2	left and right	CCA, ICA, and CB	far	automatic/semiautomatic, automatic over a specific length	end diastole	mean; maximum	higher	higher	unclear

*Insufficient information to select a specific segment or combination of segments. Place of measurement was described as follows: “the carotid bulb, internal carotid artery, and carotid bifurcation, and at least 2 cm below the bifurcation (i.e., into the common carotid artery) were imaged. CIMT was measured over a length of 1.5 cm on each carotid artery.”

CB, carotid bulb; CCA, common carotid artery; ICA, internal carotid artery; N, number; N/S, not specified.

<https://doi.org/10.1371/journal.pmed.1003414.t002>

Risk factors at the family level

Family risk factors evaluated included (1) mode of conception; and (2) pregnancy complications, such as maternal diabetes. Children conceived through ART had, on average, an increased CIMT (3 studies, 323 participants, pooled SMD 0.78 (95% CI: −0.20 to 1.75, p : 0.120), I^2 : 94%), but the imprecision around the estimate was high (Table 3; Fig 4A). There was no clear association between maternal diabetes during pregnancy and offspring's CIMT (3 studies, 658 participants, pooled SMD 0.08 (95% CI: −0.16 to 0.33, p : 0.495), I^2 : 17.1%) (Table 3; Fig 4B). Scarce evidence, reported in 1 or 2 studies, existed for other risk factors, such as preeclampsia or gestational hypertension [44,49], maternal physical activity level [42], vitamin D levels [43,68], energy and macronutrient intakes during pregnancy [42], or paternal BMI [63] (Tables 1 and S5 and S6).

Risk factors at the environmental level

Environmental risk factors evaluated included (1) exposure to tobacco (maternal, paternal, and household); and (2) socioeconomic status (maternal, family, and neighborhood). Children exposed to maternal smoking during pregnancy had, on average, an increased CIMT, but the imprecision around the estimate was high (3 studies, 909 participants, pooled SMD 0.12 (95% CI: −0.06 to 0.30, p : 0.205), I^2 : 0%) (Table 3; Fig 4C). Paternal smoking during pregnancy was linked with increased CIMT in 1 study [49]. No difference in CIMT after exposure to passive smoking in the first year of life was reported in 1 study [46]. Children with a lower socioeconomic status tended to have a higher CIMT (S5 Table). Scarce evidence, reported in 1 study,

Table 3. Summary of findings for each exposure type included in meta-analyses.

Category of exposure	Type of exposure	Level of comparison	N studies	N participants	Association measure	Association estimate	Heterogeneity
Child level							
Fetal growth	Birth size for gestational age	small* vs appropriate	16	2,570 (848 vs 1,722)	SMD	0.40 (95% CI: 0.15 to 0.64, <i>p</i> : 0.001)	<i>I</i> ² : 83%, tau ² : 0.18, <i>p</i> : <0.001
		small with IUGR diagnosis vs appropriate	8	623 (176 vs 447)	SMD	0.35 (95% CI: -0.06 to 0.77, <i>p</i> : 0.097)	<i>I</i> ² : 77%, tau ² : 0.26, <i>p</i> : <0.001
	Birth weight	range of values	7	1,445	Correlation	-0.06 (95% CI: -0.19 to 0.08, <i>p</i> : 0.404)	<i>I</i> ² : 82%, tau ² : 0.02, <i>p</i> : <0.001
	Birth length	range of values	3	424	Correlation	-0.18 (95% CI: -0.36 to 0.00, <i>p</i> : 0.052)	<i>I</i> ² : 64%, tau ² : 0.02, <i>p</i> : 0.062
	Birth head circumference	range of values	3	329	Correlation	0.01 (95% CI: -0.40 to 0.42, <i>p</i> : 0.960)	<i>I</i> ² : 93%, tau ² : 0.14, <i>p</i> : <0.001
Gestational age	Gestational age	preterm [†] vs term	7	2,024 (369 vs 1,655)	SMD	0.03 (95% CI: -0.17 to 0.22, <i>p</i> : 0.805)	<i>I</i> ² : 50%, tau ² : 0.03, <i>p</i> : 0.062
Family level							
Pregnancy-specific factors	Mode of conception	ART [‡] vs natural	3	323 (177 vs 146)	SMD	0.78 (95% CI: -0.20 to 1.75, <i>p</i> : 0.120)	<i>I</i> ² : 94%, tau ² : 0.69, <i>p</i> : <0.001
	Maternal diabetes in pregnancy	yes [§] vs no	3	658 (150 vs 508)	SMD	0.08 (95% CI: -0.16 to 0.33, <i>p</i> : 0.495)	<i>I</i> ² : 17%, tau ² : 0.01, <i>p</i> : 0.299
Environmental level							
Tobacco exposure	Maternal smoking in pregnancy	yes vs no	3	909 (145 vs 734)	SMD	0.12 (95% CI: -0.06 to 0.30, <i>p</i> : 0.205)	<i>I</i> ² : 0%, tau ² : 0, <i>p</i> : 0.380

*Birth weight and/or length below the 10th percentile or 2 SDs below the mean, with or without a documented prenatal diagnosis of IUGR (fetal biometry or Doppler velocimetry).

[†]Below 37 weeks of gestation, where specified.

[‡]In vitro fertilization or intracytoplasmic sperm injection.

[§]Gestational diabetes, where specified.

ART, assisted reproductive technologies; CI, confidence interval; IUGR, intrauterine growth restriction; *N*, number; *p*, *p*-value; SD, standard deviation; SMD, standardized mean difference; vs, versus.

<https://doi.org/10.1371/journal.pmed.1003414.t003>

existed for ambient air pollutants [57], such as nitrogen dioxide (NO₂), ozone (O₃), or particulate matter (PM₁₀ and PM_{2.5}) (Tables 1 and S6).

Discussion

Main findings


In this systematic review of 36 studies, involving 7,977 participants, multiple exposures at the child, family, or environmental levels were evaluated. Small size for gestational age had the most consistent association with increased CIMT. The magnitude of this association was higher when restricted to studies with a higher quality of CIMT measurement. The associations with conception through ART or with maternal smoking during pregnancy were not statistically significant. Evidence from interventional studies was scarce and focused exclusively on dietary interventions in the child, without showing an effect. CIMT measurement methods varied across studies, and they were frequently poorly reported, with only 33% of studies rated at higher quality across all domains of CIMT reliability.

Comparison with other studies

To the best of our knowledge, this is the first systematic review with such a multidimensional view on risk factors in the first 1,000 days of life and CIMT in children. The underlying

Small size for gestational age

(A) All



(B) Higher CIMT reliability





Fig 2. Association of small size for gestational age with CIMT in children. SMD in CIMT between children born with a small size for gestational age (exposed) and those born with an appropriate size for gestational age (reference) in (A) all studies or (B) studies at higher CIMT reliability. Weights are from random-effects model. A positive SMD corresponds to a higher CIMT in the exposed as opposed to reference. CI, confidence interval; CIMT, carotid-intima media thickness; N, sample size; p , p -value; SMD, standardized mean difference.

<https://doi.org/10.1371/journal.pmed.1003414.g002>

mechanisms of CVD programming and atherosclerosis are multifactorial. Exposure to risk factors in critical periods of development may influence life course cardiovascular health trajectories through epigenetic changes [88] and cardio-metabolic factors [89] (e.g., growth patterns,

Prematurity

(A) All



(B) Higher CIMT reliability




Fig 3. Association of prematurity with CIMT in children. SMD in CIMT between children born preterm (exposed) and those born at term (reference) in (A) all studies or (B) studies at higher CIMT reliability. Weights are from random-effects model. A positive SMD corresponds to a higher CIMT in the exposed as opposed to reference. CI, confidence interval; CIMT, carotid-intima media thickness; N, sample size; p , p -value; SMD, standardized mean difference.

<https://doi.org/10.1371/journal.pmed.1003414.g003>

blood pressure, total cholesterol, or glucose levels), but also via direct effects on the vessels structure and function. Assessing the relationship with carotid remodeling already in children was needed to shed light on the consistency and magnitude of associations, their clinical and public health importance, as well as to provide suggestions for mechanisms that need to be addressed in subsequent high-quality studies. Given that atherosclerosis is a disease affecting the whole arterial system, similar research efforts focusing on the aortic bed are currently underway (PROSPERO; CRD42019137559).

The relationship of impaired fetal growth with a higher CVD risk in later life was among the first one to be reported in the DOHaD literature. Systematic reviews and meta-analyses in adults showed that a low birth weight is associated with an increased risk of elevated blood pressure and coronary heart disease [90]. Other studies found evidence of a thicker CIMT in young adults with a small size at birth [8,9]. We found a thicker CIMT in children that were born small for gestational age, with a stronger relationship in infants than in older children. This finding is in line with previous studies and indicates that fetal growth restriction may place the individual on a higher risk trajectory since birth. It has been argued that the definition of small for gestational age may not distinguish infants that are born small due to an intra-uterine pathologic process from those that have reached their genetic potential but are constitutionally small [91]. Further, fetal growth restriction was shown to be associated with cardiac remodeling and reduced arterial compliance already in utero [92]; therefore, the severity of the growth impairment might play a key role in the vascular abnormalities of these children. Following this reasoning, we restricted our analyses to children that were born small for

Table 4. Subgroup meta-analyses of SMD for the association of small size for gestational age or prematurity with CIMT.

	Birth size for gestational age (small vs appropriate)				Gestational age (preterm vs term)			
	N studies	SMD	I ² , tau ² *	p [†]	N studies	SMD	I ² , tau ² *	p [†]
All	16	0.40 (95% CI: 0.15 to 0.64, <i>p</i> : 0.001)	83%, 0.18	-	7	0.03 (95% CI: -0.17 to 0.22, <i>p</i> : 0.805)	50%, 0.03	-
Study design								
Cohort	7	0.39 (95% CI: 0 to 0.78, <i>p</i> : 0.049)	79%, 0.2	0.528	5	0.1 (95% CI: -0.13 to 0.33, <i>p</i> : 0.383)	45%, 0.03	0.030
Cross-sectional	9	0.41 (95% CI: 0.07 to 0.74, <i>p</i> : 0.017)	87%, 0.21		2	-0.18 (95% CI: -0.39 to 0.04, <i>p</i> : 0.104)	0%, 0	
Study setting								
Community or population based	1	-0.09 (95% CI: -0.30 to 0.13, <i>p</i> : 0.440)	-	<0.001	3	0.08 (95% CI: -0.23 to 0.38, <i>p</i> : 0.626)	73%, 0.05	0.716
Healthcare facility	13	0.43 (95% CI: 0.14 to 0.72, <i>p</i> : 0.004)	83%, 0.21		4	-0.04 (95% CI: -0.34 to 0.27, <i>p</i> : 0.816)	32%, 0.03	
N/S	2	0.60 (95% CI: 0.36 to 0.85, <i>p</i> : <0.001)	1%, 0.00		-	-	-	
CIMT wall								
Far	11	0.38 (95% CI: 0.11 to 0.64, <i>p</i> : 0.005)	82%, 0.15	0.756	6	0.08 (95% CI: -0.11 to 0.27, <i>p</i> : 0.425)	42%, 0.02	0.066
N/S	5	0.47 (95% CI: -0.17 to 1.12, <i>p</i> : 0.149)	88%, 0.46		1	-0.37 (95% CI: -0.81 to 0.06, <i>p</i> : 0.090)	-	
CIMT sides								
Left and right	12	0.5 (95% CI: 0.22 to 0.79, <i>p</i> : 0.001)	84%, 0.19	0.196	4	0.02 (95% CI: -0.2 to 0.23, <i>p</i> : 0.889)	41%, 0.02	0.059
Right	2	0 (95% CI: -0.35 to 0.35, <i>p</i> : 1.000)	0%, 0.00		1	0.35 (95% CI: 0.01 to 0.69, <i>p</i> : 0.045)	-	
N/S	2	0.11 (95% CI: -1.00 to 1.21, <i>p</i> : 0.849)	95%, 0.60		2	-0.22 (95% CI: -0.6 to 0.17, <i>p</i> : 0.272)	19%, 0.02	
CIMT edge detection method								
Automatic or semiautomatic	4	0.35 (95% CI: 0.04 to 0.67, <i>p</i> : 0.029)	69%, 0.06	<0.001	3	0.21 (95% CI: 0.03 to 0.39, <i>p</i> : 0.021)	0%, 0	0.006
Manual	6	0.22 (95% CI: -0.26 to 0.71, <i>p</i> : 0.367)	88%, 0.31		2	-0.23 (95% CI: -0.43 to -0.03, <i>p</i> : 0.025)	0%, 0	
N/S	6	0.64 (95% CI: 0.27 to 1.01, <i>p</i> : 0.001)	71%, 0.14		2	-0.02 (95% CI: -0.46 to 0.43, <i>p</i> : 0.949)	0%, 0	
Ultrasound transducer frequency								
≤12 MHz	10	0.69 (95% CI: 0.45 to 0.93, <i>p</i> : <0.001)	57%, 0.07	<0.001	3	0.21 (95% CI: -0.06 to 0.48, <i>p</i> : 0.123)	0%, 0	0.253
>12 MHz	4	0.03 (95% CI: -0.5 to 0.56, <i>p</i> : 0.904)	90%, 0.26		2	-0.06 (95% CI: -0.66 to 0.54, <i>p</i> : 0.846)	77%, 0.14	
N/S	2	-0.06 (95% CI: -0.25 to 0.12, <i>p</i> : 0.505)	0%, 0.00		2	-0.04 (95% CI: -0.34 to 0.27, <i>p</i> : 0.808)	69%, 0.03	
Age								
0 to 1 years	4	0.63 (95% CI: -0.02 to 1.27, <i>p</i> : 0.057)	91%, 0.39	0.002	1	0.03 (95% CI: -0.53 to 0.59, <i>p</i> : 0.924)	-	0.957
2 to 16 years	12	0.31 (95% CI: 0.06 to 0.55, <i>p</i> : 0.014)	77%, 0.13		6	0.02 (95% CI: -0.2 to 0.25, <i>p</i> : 0.828)	58%, 0.04	
N participants								
<200	11	0.44 (95% CI: 0.05 to 0.84, <i>p</i> : 0.027)	83%, 0.35	0.699	4	-0.04 (95% CI: -0.34 to 0.27, <i>p</i> : 0.816)	32%, 0.03	0.716
≥200	5	0.34 (95% CI: 0.03 to 0.65, <i>p</i> : 0.034)	86%, 0.11		3	0.08 (95% CI: -0.23 to 0.38, <i>p</i> : 0.626)	73%, 0.05	

*I² and tau² assess within-group heterogeneity.†*p*-value assesses between-group heterogeneity.CI, confidence interval; CIMT, carotid intima-media thickness; N, number; N/S, not specified; *p*, *p*-value; SMD, standardized mean difference.<https://doi.org/10.1371/journal.pmed.1003414.t004>




(A) Assisted reproductive technology conception**(B) Maternal diabetes in pregnancy****(C) Maternal smoking in pregnancy**

Fig 4. Association of (A) ART conception, (B) maternal diabetes in pregnancy, and (C) maternal smoking in pregnancy with CIMT in children. SMD in CIMT between children: (A) conceived through ART (exposed) or naturally (reference); (B) exposed to maternal diabetes during pregnancy (exposed) or not exposed (reference); and (C) exposed to maternal smoking during pregnancy (exposed) or not exposed (reference). Weights are from random-effects model. A positive SMD corresponds to a higher CIMT in the exposed as opposed to reference. ART, assisted reproductive technologies; CI, confidence interval; CIMT, carotid-intima media thickness; N, sample size; p , p -value; SMD, standardized mean difference.

<https://doi.org/10.1371/journal.pmed.1003414.g004>

gestational age and had a documented prenatal diagnosis of IUGR. CIMT remained thicker in IUGR children compared with those born appropriate for gestational age (Table 3). Also, the vast majority of included studies used Doppler velocimetry in the maternal or fetal vessels to diagnose IUGR; thus, these findings support the role of a well-characterized mechanism of vascular remodeling after poor fetal growth, namely the placental dysfunction [10].

The duration of gestation, maternal diabetes, or parental smoking during pregnancy are known determinants of the size at birth [1] and may also directly affect the vascular function and structure. Prematurity was shown to increase the risk of hypertension in adolescents and adults [93], probably through impaired renal development. Young adults that were born pre-term were shown to have a slightly higher CIMT, noted in 3 of 4 studies of a recent systematic review and meta-analysis [94]. We did not find consistent evidence of a relationship between prematurity and higher CIMT in children. This may be because prematurity is more likely to be associated with a higher cardiovascular risk only if accompanied by fetal growth restriction or higher blood pressure in later life [8,95]. Also, it may be due to the measurement error in CIMT. In our case, it is plausible to be a combination of these mechanisms, as studies meta-

analyzed involved former preterm children, with or without poor fetal growth; therefore, a distortion of this association cannot be excluded, and CIMT was higher when restricted to studies with more reliable measurements. Of note, the relationship of CIMT with prematurity was weaker than that with small for gestational age, which pleads for a greater impact of the latter, and potentially, a more pronounced impact when the 2 risk factors are present concomitantly. Interestingly, we found inverse, but very weak, relationships of CIMT with other growth indicators, such as birth length, potentially because they do not account for the duration of gestation. We also found no clear evidence of a higher CIMT after exposure to maternal diabetes during pregnancy and a consistently higher CIMT, although of very low magnitude, in children exposed to maternal smoking during pregnancy. It is known that gestational diabetes is associated with macrosomia in the offspring, whereas maternal smoking during pregnancy is associated with poor fetal growth [1]. Our data indicate that CIMT was consistently increased in children with abnormal growth, either large or small. The effect of maternal gestational diabetes or smoking may thus be mainly indirect, through the size at birth. However, adequately powered studies that formally assess the mediator effect of size at birth on these relationships are needed to shed light on this hypothesis [96]. Finally, our finding of a higher CIMT in children conceived through ART draws attention on the highly sensitive periconceptional period, with potentially new underlying mechanisms that need to be disentangled.

Heterogeneity and poorly reported or missing information on key characteristics of the CIMT measurement methods were among the main issues that we encountered when analyzing the evidence presented herein. Similar conclusions were drawn in systematic reviews on CIMT in adults that carefully considered measurement methods when interpreting their results [6,97]. In our case, the sensitivity analyses performed for small size for gestational age and prematurity showed that the quality of CIMT measurement methods had a significant impact on the results. Additionally, CIMT values may differ with the edge detection method and the specificities of the ultrasound equipment used, such as the transducer frequency. We could notice large decreases in the magnitude of association estimates across studies using higher transducer frequencies. Our finding is in line with other evidence showing that conventional high frequencies (≤ 12 MHz) may have insufficient ultrasound resolution and result in an overestimation of the true thickness in children under 12 years of age [18,19]. Novel techniques that measure CIMT with very high-resolution ultrasound (25 to 55 MHz) seem to be more precise in infants and young children and at very early stages of atherosclerosis [98,99].

Strengths and limitations

The strength of our systematic review lies primarily in the fact that it was performed according to a detailed protocol that was prospectively registered on PROSPERO and published in full in a peer-reviewed journal. Secondly, we collected extensive data on the CIMT measurement characteristics, which permitted interpretation of results in the light of the quality of measurement. Thirdly, we included both observational and interventional studies reporting on exposures or interventions at the child, family, or environmental levels. This helped us have a comprehensive overview on the current status of the evidence and highlight needed areas of future research.

The main limitation is that the body of evidence for this systematic review is largely observational, which is of low certainty by design and restricts the ability to draw conclusions about causality. Additionally, the degree of confidence in results is limited by the high between-study heterogeneity. Studies varied on several aspects, such as design, exposure metric, age at CIMT assessment, or CIMT image analysis. The variability in age at CIMT assessment was large, from 0 to 18 years. Our subgroup analysis was mainly based on the age range reported in each

study and, thus, could only distinguish the infants from the other pediatric age groups combined. Meta-analyses were performed mainly using estimates from bivariate associations. Data transformation and simplification were needed, and associations were estimated from various statistics so that relationships could be summarized in a consistent way across studies. Therefore, differences in body size between exposed and nonexposed could not be taken into account in the analyses, confounding bias cannot be excluded, and our association estimates may be distorted, probably underestimated, due to measurement error in the CIMT. Furthermore, CIMT was evaluated in youth with tiny, submillimetric vessel structures, and many studies had relatively small samples and few individuals exposed to the risk factors of interest, which resulted in large CIs for the estimates. The low number of studies meta-analyzed often impeded conduction of subgroup analyses to explore the sources of heterogeneity predefined in the protocol. However, to increase our confidence in the overall estimates, whenever possible, we conducted analyses restricted to studies at higher quality for all CIMT reliability domains. Other limitations are represented by exclusion of studies not reported in English or French [100–102], publication bias, and selective exposure or analysis reporting. Usable data for the majority of exposures or interventions were reported by a low to a very low number of studies. However, for small size for gestational age, the risk factor most studied, the funnel plot inspection, and Egger test were not indicative of publication bias (S2 Fig).

Implications for research and practice

We highlight the need of a standardized ultrasound protocol for measuring CIMT in children. Due to the paucity of interventional studies and the limited quality of CIMT measurements, further high-quality studies are needed to justify the use of CIMT for child CVD risk assessment in clinical practice. However, promotion of a healthy lifestyle is important at any age, and screening of postnatal CVD factors, such as hypertension or obesity, targeted at children exposed to risk factors in the first 1,000 days of life, may be warranted.

Further, our study indicated that children that were small for gestational age had a higher CIMT in early life. Prevention of small for gestational age could have a big impact from a public health perspective. Its prevalence varies among populations and rises with decreasing gestational age and availability of resources, reaching about 20% in low and middle-income countries [103]. Moreover, modifiable factors, including gestational smoking or poor maternal nutrition, account for a large share of small for gestational age infants [104]. In fact, smoking cessation interventions during pregnancy were shown to result in a 17% reduction in low birth weight cases [105]. Public health strategies to reduce smoking and improve the nutritional status in women of reproductive age should be advocated and may improve the cardiovascular health too [105,106].

Conclusions

In our meta-analyses, we found several risk factors in the first 1,000 days of life that may be associated with increased CIMT in infants, children, and adolescents. Poor fetal growth had the most consistent and strongest association with increased CIMT. The associations with ART conception or with maternal smoking during pregnancy were not statistically significant. From a DOHaD perspective, exposure to adverse experiences in early life determines adaptive responses in the organs' structure and function with lifelong effects on cardiovascular health and disease, ranging from elevated blood pressure, early vascular aging and increased CIMT, and premature CVD morbidity and mortality [1,107]. Therefore, assessing CIMT early in life is important to distinguish between changes occurring in childhood and adulthood, with the goal of better characterizing lifetime risk trajectories. Provided this hypothesis will stand the test of time and effective intervention, the interplay of risk factors in the first 1,000

days of life and vascular remodeling will offer a great opportunity for primordial prevention of CVD [108].

Supporting information

S1 PRISMA Checklist. Preferred Reporting Items for Systematic Reviews and Meta-Analyses.

(PDF)

S1 Fig. Assessment of study quality for each exposure type included in meta-analyses.

(PDF)

S2 Fig. Assessment of small-study effects, including publication bias, for each exposure type included in meta-analyses.

(PDF)

S3 Fig. Random-effects meta-regression examining the influence of sample size on the association of small size for gestational age with CIMT.

(PDF)

S1 Table. Strategies for systematic searches.

(PDF)

S2 Table. Strategies for supplementary searches.

(PDF)

S3 Table. Criteria for CIMT quality assessment.

(PDF)

S4 Table. CIMT equipment and operator.

(PDF)

S5 Table. Association of CIMT with main exposures or interventions types in the first 1,000 days of life.

(PDF)

S6 Table. Other exposure types in the first 1,000 days of life by level and category of exposure (reported in a single study).

(PDF)

S7 Table. Criteria for study quality assessment.

(PDF)

S8 Table. Assessment of study quality for each intervention type in interventional studies.

(PDF)

S9 Table. Assessment of study quality for observational studies included in meta-analyses.

(PDF)

Acknowledgments

The authors thank Thomas Brauchli (Data and Documentation Unit, Center for Primary Care and Public Health (UNISANTÉ), University of Lausanne, Lausanne, Switzerland) and Cécile Jaques (Medical Library, Lausanne University Hospital, University of Lausanne, Lausanne, Switzerland), librarians, who assisted with the development of the literature search strategies and provided technical advice for running the search queries in the databases.

Author Contributions

Conceptualization: Adina Mihaela Epure, Arnaud Chiolerio, Nicole Sekarski.

Data curation: Adina Mihaela Epure.

Formal analysis: Adina Mihaela Epure, Magali Rios-Leyvraz, Daniela Anker, Bruno R. da Costa, Arnaud Chiolerio.

Funding acquisition: Arnaud Chiolerio.

Investigation: Adina Mihaela Epure, Magali Rios-Leyvraz, Daniela Anker, Stefano Di Bernardo, Arnaud Chiolerio, Nicole Sekarski.

Methodology: Adina Mihaela Epure, Stefano Di Bernardo, Bruno R. da Costa, Arnaud Chiolerio, Nicole Sekarski.

Project administration: Adina Mihaela Epure, Arnaud Chiolerio.

Supervision: Bruno R. da Costa, Arnaud Chiolerio, Nicole Sekarski.

Visualization: Adina Mihaela Epure.

Writing – original draft: Adina Mihaela Epure.

Writing – review & editing: Adina Mihaela Epure, Magali Rios-Leyvraz, Daniela Anker, Stefano Di Bernardo, Bruno R. da Costa, Arnaud Chiolerio, Nicole Sekarski.

References

1. Hanson MA, Gluckman PD. Early developmental conditioning of later health and disease: physiology or pathophysiology? *Physiol Rev.* 2014; 94(4):1027–1076. <https://doi.org/10.1152/physrev.00029.2013> PMID: 25287859
2. Fleming TP, Watkins AJ, Velazquez MA, Mathers JC, Prentice AM, Stephenson J, et al. Origins of life-time health around the time of conception: causes and consequences. *Lancet.* 2018; 391(10132):1842–1852. [https://doi.org/10.1016/S0140-6736\(18\)30312-X](https://doi.org/10.1016/S0140-6736(18)30312-X) PMID: 29673874
3. Huxley RR, Shiell AW, Law CM. The role of size at birth and postnatal catch-up growth in determining systolic blood pressure: a systematic review of the literature. *J Hypertens.* 2000; 18(7):815–831. <https://doi.org/10.1097/00004872-200018070-00002> PMID: 10930178
4. Tian G, Guo C, Li Q, Liu Y, Sun X, Yin Z, et al. Birth weight and risk of type 2 diabetes: A dose-response meta-analysis of cohort studies. *Diabetes Metab Res Rev.* 2019; 35(5):e3144. <https://doi.org/10.1002/dmrr.3144> PMID: 30786122
5. Risnes KR, Vatten LJ, Baker JL, Jameson K, Sovio U, Kajantie E, et al. Birthweight and mortality in adulthood: a systematic review and meta-analysis. *Int J Epidemiol.* 2011; 40(3):647–661. <https://doi.org/10.1093/ije/dyq267> PMID: 21324938
6. Lorenz MW, Markus HS, Bots ML, Rosvall M, Sitzer M. Prediction of clinical cardiovascular events with carotid intima-media thickness: a systematic review and meta-analysis. *Circulation.* 2007; 115(4):459–467. <https://doi.org/10.1161/CIRCULATIONAHA.106.628875> PMID: 17242284
7. Eikendal AL, Groenewegen KA, Anderson TJ, Britton AR, Engstrom G, Evans GW, et al. Common carotid intima-media thickness relates to cardiovascular events in adults aged <45 years. *Hypertension.* 2015; 65(4):707–713. <https://doi.org/10.1161/HYPERTENSIONAHA.114.04658> PMID: 25624341
8. Skilton MR, Viikari JS, Juonala M, Laitinen T, Lehtimäki T, Taittonen L, et al. Fetal growth and preterm birth influence cardiovascular risk factors and arterial health in young adults: the Cardiovascular Risk in Young Finns Study. *Arterioscler Thromb Vasc Biol.* 2011; 31(12):2975–2981. <https://doi.org/10.1161/ATVBAHA.111.234757> PMID: 21940950
9. Oren A, Vos LE, Uiterwaal CS, Gorissen WH, Grobbee DE, Bots ML. Birth weight and carotid intima-media thickness: new perspectives from the atherosclerosis risk in young adults (ARYA) study. *Ann Epidemiol.* 2004; 14(1):8–16. [https://doi.org/10.1016/s1047-2797\(03\)00068-1](https://doi.org/10.1016/s1047-2797(03)00068-1) PMID: 14664774
10. Visentin S, Grumolato F, Nardelli GB, Di Camillo B, Grisan E, Cosmi E. Early origins of adult disease: low birth weight and vascular remodeling. *Atherosclerosis.* 2014; 237(2):391–399. <https://doi.org/10.1016/j.atherosclerosis.2014.09.027> PMID: 25463063

11. Crispi F, Bijlens B, Figueras F, Bartrons J, Eixarch E, Le Noble F, et al. Fetal growth restriction results in remodeled and less efficient hearts in children. *Circulation*. 2010; 121(22):2427–2436. <https://doi.org/10.1161/CIRCULATIONAHA.110.937995> PMID: 20497977
12. Dratva J, Breton CV, Hodis HN, Mac KWJ, Salam MT, Zemp E, et al. Birth weight and carotid artery intima-media thickness. *J Pediatr*. 2013; 162(5):906–11.e2. <https://doi.org/10.1016/j.jpeds.2012.10.060> PMID: 23260106
13. Jouret B, Dulac Y, Bassil Eter R, Taktak A, Cristini C, Lounis N, et al. Endothelial function and mechanical arterial properties in children born small for gestational age: comparison with obese children. *Horm Res Paediatr*. 2011; 76(4):240–247. <https://doi.org/10.1159/000329379> PMID: 21912079
14. Stock K, Schmid A, Griesmaier E, Gande N, Hochmayr C, Knoflach M, et al. The Impact of Being Born Preterm or Small for Gestational Age on Early Vascular Aging in Adolescents. *J Pediatr*. 2018; 201:49–54.e1. <https://doi.org/10.1016/j.jpeds.2018.05.056> PMID: 29960764
15. Hanson M, Gluckman P. Developmental origins of noncommunicable disease: population and public health implications. *Am J Clin Nutr*. 2011; 94(6 Suppl):1754s–1758s. <https://doi.org/10.3945/ajcn.110.001206> PMID: 21525196
16. Dalla Pozza R, Ehringer-Schetitska D, Fritsch P, Jokinen E, Petropoulos A, Oberhoffer R. Intima media thickness measurement in children: A statement from the Association for European Paediatric Cardiology (AEPC) Working Group on Cardiovascular Prevention endorsed by the Association for European Paediatric Cardiology. *Atherosclerosis*. 2015; 238(2):380–387. <https://doi.org/10.1016/j.atherosclerosis.2014.12.029> PMID: 25555270
17. Urbina EM, Williams RV, Alpert BS, Collins RT, Daniels SR, Hayman L, et al. Noninvasive assessment of subclinical atherosclerosis in children and adolescents: recommendations for standard assessment for clinical research: a scientific statement from the American Heart Association. *Hypertension*. 2009; 54(5):919–950. <https://doi.org/10.1161/HYPERTENSIONAHA.109.192639> PMID: 19729599
18. Sarkola T, Redington A, Keeley F, Bradley T, Jaeggi E. Transcutaneous very-high-resolution ultrasound to quantify arterial wall layers of muscular and elastic arteries: validation of a method. *Atherosclerosis*. 2010; 212(2):516–523. <https://doi.org/10.1016/j.atherosclerosis.2010.06.043> PMID: 20673898
19. Sarkola T, Slorach C, Hui W, Bradley TJ, Redington AN, Jaeggi E. Transcutaneous very-high resolution ultrasound for the quantification of carotid arterial intima-media thickness in children—feasibility and comparison with conventional high resolution vascular ultrasound imaging. *Atherosclerosis*. 2012; 224(1):102–107. <https://doi.org/10.1016/j.atherosclerosis.2012.06.054> PMID: 22784638
20. Potter K, Reed CJ, Green DJ, Hankey GJ, Arnolda LF. Ultrasound settings significantly alter arterial lumen and wall thickness measurements. *Cardiovasc Ultrasound*. 2008; 6:6. <https://doi.org/10.1186/1476-7120-6-6> PMID: 18208623
21. de Groot E, Hovingh GK, Wiegman A, Duriez P, Smit AJ, Fruchart JC, et al. Measurement of arterial wall thickness as a surrogate marker for atherosclerosis. *Circulation*. 2004; 109(23 Suppl 1):Ii33–Ii38. <https://doi.org/10.1161/01.CIR.0000131516.65699.ba> PMID: 15198964
22. Mivelaz Y, Di Bernardo S, Boulou Ksontini T, Prsa M, Vial Y, Chiolerio A, et al. Feasibility and reliability of carotid intima-media thickness measurements in nonsedated infants. *J Hypertens*. 2016; 34(11):2227–2232. <https://doi.org/10.1097/HJH.0000000000001065> PMID: 27490951
23. Epure AM, Leyvraz M, Mivelaz Y, Di Bernardo S, da Costa BR, Chiolerio A, et al. Risk factors and determinants of carotid intima-media thickness in children: protocol for a systematic review and meta-analysis. *BMJ Open*. 2018; 8(6):e019644. <https://doi.org/10.1136/bmjopen-2017-019644> PMID: 29866720
24. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gotzsche PC, Ioannidis JP, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ*. 2009; 339:b2700. <https://doi.org/10.1136/bmj.b2700> PMID: 19622552
25. Stroup DF, Berlin JA, Morton SC, Olkin I, Williamson GD, Rennie D, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *JAMA*. 2000; 283(15):2008–2012. <https://doi.org/10.1001/jama.283.15.2008> PMID: 10789670
26. Moore TG, Arefadib N, Deery A, Keyes M, West S. The first thousand days: an evidence paper—summary. Parkville, Victoria: Centre for Community Child Health, Murdoch Children's Research Institute; 2017.
27. Bramer WM, Giustini D, de Jonge GB, Holland L, Bekhuis T. De-duplication of database search results for systematic reviews in EndNote. *J Med Libr Assoc*. 2016; 104(3):240–243. <https://doi.org/10.3163/1536-5050.104.3.014> PMID: 27366130

28. Wells G, Shea B, O'Connell D, Peterson J, Welch V, Losos M, et al. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. [cited 2017 Aug]. Available from: http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp.
29. Higgins JPT, Green S. Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0: The Cochrane Collaboration. 2011. Available from: <http://www.handbook-5-1.cochrane.org>.
30. Higgins JP, Altman DG, Gøtzsche PC, Jüni P, Moher D, Oxman AD, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ*. 2011; 343:d5928. <https://doi.org/10.1136/bmj.d5928> PMID: 22008217
31. Cooper H, Hedges LV, Valentine JC, editors. *The Handbook of Research Synthesis and Meta-Analysis*. Russell Sage Foundation; 2009.
32. Borenstein M, Hedges LV, Higgins JPT, Rothstein HR. *Introduction to Meta-Analysis*. John Wiley & Sons; 2009. <https://doi.org/10.1002/9780470743386> PMID: 29218357
33. Lipsey MW, Wilson DB. *Practical Meta-Analysis*. SAGE Publications Inc; 2000.
34. Rücker G, Cates CJ, Schwarzer G. Methods for including information from multi-arm trials in pairwise meta-analysis. *Res Synth Methods*. 2017; 8(4):392–403. <https://doi.org/10.1002/jrsm.1259> PMID: 28759708
35. Wan X, Wang W, Liu J, Tong T. Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range. *BMC Med Res Methodol*. 2014; 14:135. <https://doi.org/10.1186/1471-2288-14-135> PMID: 25524443
36. Aloe AM, Thompson CG. The synthesis of partial effect sizes. *J Soc Soc Work Res*. 2013; 4(4):390–405. <https://doi.org/10.5243/jsswr.2013.24>
37. Nieminen P, Lehtiniemi H, Vähäkangas K, Huusko A, Rautio A. Standardised regression coefficient as an effect size index in summarising findings in epidemiological studies. *Epidemiol Biostatist Public Health*. 2013; 10. <https://doi.org/10.2427/8854>
38. Lüdtke D. *esc: Effect Size Computation for Meta Analysis (Version 0.5.1)*. 2019. <https://doi.org/10.5281/zenodo.1249218> Available from: <https://CRAN.R-project.org/package=esc>.
39. Aloe AM. Inaccuracy of regression results in replacing bivariate correlations. *Res Synth Methods*. 2015; 6(1):21–27. <https://doi.org/10.1002/jrsm.1126> PMID: 26035467
40. Amrhein V, Greenland S, McShane B. Scientists rise up against statistical significance. *Nature*. 2019; 567(7748):305–307. <https://doi.org/10.1038/d41586-019-00857-9> PMID: 30894741
41. Higgins JPT, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ*. 2003; 327(7414):557–560. <https://doi.org/10.1136/bmj.327.7414.557> PMID: 12958120
42. Gale CR, Jiang B, Robinson SM, Godfrey KM, Law CM, Martyn CN. Maternal diet during pregnancy and carotid intima-media thickness in children. *Arterioscler Thromb Vasc Biol*. 2006; 26(8):1877–1882. <https://doi.org/10.1161/01.ATV.0000228819.13039.b8> PMID: 16728657
43. Gale CR, Robinson SM, Harvey NC, Javaid MK, Jiang B, Martyn CN, et al. Maternal vitamin D status during pregnancy and child outcomes. *Eur J Clin Nutr*. 2008; 62(1):68–77. <https://doi.org/10.1038/sj.ejcn.1602680> PMID: 17311057
44. Ayer JG, Harmer JA, Nakhla S, Xuan W, Ng MKC, Raitakari OT, et al. HDL-cholesterol, blood pressure, and asymmetric dimethylarginine are significantly associated with arterial wall thickness in children. *Arterioscler Thromb Vasc Biol*. 2009a; 29(6):943–949. <https://doi.org/10.1161/ATVBAHA.109.184184> PMID: 19359663
45. Ayer JG, Harmer JA, Xuan W, Toelle B, Webb K, Almqvist C, et al. Dietary supplementation with n-3 polyunsaturated fatty acids in early childhood: Effects on blood pressure and arterial structure and function at age 8 y. *Am J Clin Nutr*. 2009b; 90(2):438–446. <https://doi.org/10.3945/ajcn.2009.27811> PMID: 19515739
46. Ayer JG, Belousova E, Harmer JA, David C, Marks GB, Celermajor DS. Maternal cigarette smoking is associated with reduced high-density lipoprotein cholesterol in healthy 8-year-old children. *Eur Heart J*. 2011; 32(19):2446–2453. <https://doi.org/10.1093/eurheartj/ehr174> PMID: 21693475
47. Skilton MR, Ayer JG, Harmer JA, Webb K, Leeder SR, Marks GB, et al. Impaired fetal growth and arterial wall thickening: A randomized trial of omega-3 supplementation. *Pediatrics*. 2012; 129(3):e698–e703. <https://doi.org/10.1542/peds.2011-2472> PMID: 22351892
48. Skilton MR, Marks GB, Ayer JG, Garden FL, Garnett SP, Harmer JA, et al. Weight gain in infancy and vascular risk factors in later childhood. *Pediatrics*. 2013; 131(6):e1821–e1828. <https://doi.org/10.1542/peds.2012-2789> PMID: 23713097
49. Rodriguez-Lopez M, Osorio L, Acosta-Rojas R, Figueras J, Cruz-Lemini M, Figueras F, et al. Influence of breastfeeding and postnatal nutrition on cardiovascular remodeling induced by fetal growth restriction. *Pediatr Res*. 2016; 79(1):100–106. <https://doi.org/10.1038/pr.2015.182> PMID: 26372518

50. Trevisanuto D, Avezú F, Cavallin F, Doglioni N, Marzolo M, Verlato F, et al. Arterial wall thickness and blood pressure in children who were born small for gestational age: Correlation with umbilical cord high-sensitivity C-reactive protein. *Arch Dis Child*. 2010; 95(1):31–34. <https://doi.org/10.1136/adc.2008.150326> PMID: 19773221
51. Evelein AMV, Geerts CC, Visseren FLJ, Bots ML, Van Der Ent CK, Grobbee DE, et al. The association between breastfeeding and the cardiovascular system in early childhood. *Am J Clin Nutr*. 2011; 93(4):712–718. <https://doi.org/10.3945/ajcn.110.002980> PMID: 21310835
52. Geerts CC, Bots ML, van der Ent CK, Grobbee DE, Uiterwaal CSPM. Parental smoking and vascular damage in their 5-year-old children. *Pediatrics*. 2012; 129(1):45–54. <https://doi.org/10.1542/peds.2011-0249> PMID: 22201150
53. Evelein AMV, Visseren FLJ, Van Der Ent CK, Grobbee DE, Uiterwaal CSPM. Excess early postnatal weight gain leads to thicker and stiffer arteries in young children. *J Clin Endocrinol Metab*. 2013; 98(2):794–801. <https://doi.org/10.1210/jc.2012-3208> PMID: 23284005
54. Pluymen LPM, Dalmeijer GW, Smit HA, Uiterwaal C, van der Ent CK, van Rossem L. Long-chain polyunsaturated fatty acids in infant formula and cardiovascular markers in childhood. *Matern Child Nutr*. 2017; 14(2):e12523. <https://doi.org/10.1111/mcn.12523> PMID: 28949070
55. Atabek ME, Çaan HH, Ekliolu BS, Oran B. Absence of increase in carotid artery Intima-Media thickness in infants of diabetic mothers. *J Clin Res Pediatr Endocrinol*. 2011; 3(3):144–148. <https://doi.org/10.4274/jcrpe.v3i3.28> PMID: 21911328
56. Breton CV, Gao L, Yao J, Siegmund KD, Lurmann F, Gilliland F. Particulate matter, the newborn methylome, and cardio-respiratory health outcomes in childhood. *Environ Epigenet*. 2016b; 2(2):dvw005. <https://doi.org/10.1093/eep/dvw005> PMID: 29492287
57. Breton CV, Yao J, Millstein J, Gao L, Siegmund KD, Mack W, et al. Prenatal air pollution exposures, DNA methyl transferase genotypes, and associations with newborn line1 and Alu methylation and childhood blood pressure and carotid intima-media thickness in the children's health study. *Environ Health Perspect*. 2016a; 124(12):1905–1912. <https://doi.org/10.1289/EHP181> PMID: 27219456
58. Schubert U, Müller M, Abdul-Khaliq H, Norman M, Bonamy AKE. Relative intima-media thickening after preterm birth. *Acta Paediatr*. 2013; 102(10):965–969. <https://doi.org/10.1111/apa.12355> PMID: 23848508
59. Valenzuela-Alcaraz B, Serafini A, Sepulveda-Martinez A, Casals G, Rodriguez-Lopez M, Garcia-Otero L, et al. Postnatal persistence of fetal cardiovascular remodelling associated with assisted reproductive technologies: a cohort study. *BJOG*. 2019; 126(2):291–298. <https://doi.org/10.1111/1471-0528.15246> PMID: 29673050
60. Lee H, Dichtl S, Mormanova Z, Dalla Pozza R, Genzel-Boroviczeny O. In adolescence, extreme prematurity is associated with significant changes in the microvasculature, elevated blood pressure and increased carotid intima—Media thickness. *Arch Dis Child Educ Pract Ed*. 2014; 99(10):907–911. <https://doi.org/10.1136/archdischild-2013-304074> PMID: 24879769
61. Morsing E, Liuba P, Fellman V, Maršál K, Brodzki J. Cardiovascular function in children born very pre-term after intrauterine growth restriction with severely abnormal umbilical artery blood flow. *Eur J Prev Cardiol*. 2014; 21(10):1257–1266. <https://doi.org/10.1177/2047487313486044> PMID: 23613223
62. Stergiotou I, Crispi F, Valenzuela-Alcaraz B, Cruz-Lemini M, Bijns B, Gratacos E. Aortic and carotid intima-media thickness in term small-for-gestational-age newborns and relationship with prenatal signs of severity. *Ultrasound Obstet Gynecol*. 2014; 43(6):625–631. <https://doi.org/10.1002/uog.13245> PMID: 24272754
63. Gruszfeld D, Weber M, Nowakowska-Rysz M, Janas R, Kozlik-Feldmann R, Khonneux A, et al. Protein intake in infancy and carotid intima media thickness at 5 years—A secondary analysis from a randomized trial for the European childhood obesity study group. *Ann Nutr Metab*. 2015; 66(1):51–59. <https://doi.org/10.1159/000369980> PMID: 25572773
64. Sebastiani G, Díaz M, Bassols J, Aragonés G, López-Bermejo A, de Zegher F, et al. The sequence of prenatal growth restraint and post-natal catch-up growth leads to a thicker intima-media and more preperitoneal and hepatic fat by age 3–6 years. *Pediatr Obes*. 2016; 11(4):251–257. <https://doi.org/10.1111/ijpo.12053> PMID: 26132470
65. Liu RS, Mensah FK, Carlin J, Edwards B, Ranganathan S, Cheung M, et al. Socioeconomic Position Is Associated With Carotid Intima-Media Thickness in Mid-Childhood: The Longitudinal Study of Australian Children. *J Am Heart Assoc*. 2017; 6(8). <https://doi.org/10.1161/jaha.117.005925> PMID: 28862928
66. Mohlkert LA, Hallberg J, Broberg O, Hellström M, Pegelow Halvorsen C, Sjöberg G, et al. Preterm arteries in childhood: Dimensions, intima-media thickness, and elasticity of the aorta, coronaries, and carotids in 6-y-old children born extremely preterm. *Pediatr Res*. 2017; 81(2):299–306. <https://doi.org/10.1038/pr.2016.212> PMID: 28195589

67. Tzschoppe A, Von Kries R, Struwe E, Rascher W, Dörr HG, Jüngert J, et al. Intrauterine Growth Restriction (IUGR) Induces Signs of Subclinical Atherosclerosis in 6-Year-Old Infants Despite Absence of Excessive Growth. *Klinische Padiatrie*. 2017; 229(4):209–215. <https://doi.org/10.1055/s-0043-104528> PMID: 28718186
68. Carreras-Badosa G, Armero-Bujaldón C, Solé-Amat L, Prats-Puig A, Díaz-Roldán F, Soriano-Rodriguez P, et al. Serum 25-hydroxyvitamin D and cardiovascular disease risk factors in women with excessive weight gain during pregnancy and in their offspring at age 5–6 years. *Int J Obes*. 2018; 42(5):1019–1028. <https://doi.org/10.1038/s41366-018-0101-6> PMID: 29777240
69. Chen L, Guilmette J, Luo ZC, Cloutier A, Wang WJ, Yang MN, et al. Placental 11beta-HSD2 and Cardiometabolic Health Indicators in Infancy. *Diabetes Care*. 2019. <https://doi.org/10.2337/dc18-2041> PMID: 30833369
70. Prins-Van Ginkel AC, Bruijning-Verhagen PCJ, Wijga AH, Bots ML, Gehring U, Van Der Hoek W, et al. Childhood infections and common carotid intima media thickness in adolescence. *Epidemiol Infect*. 2019; 147. <https://doi.org/10.1017/S095026881800287X> PMID: 30404679
71. Sebastiani G, García-Beltrán C, Pie S, Guerra A, López-Bermejo A, de Toledo JS, et al. The sequence of prenatal growth restraint and postnatal catch-up growth: normal heart but thicker intima-media and more pre-peritoneal fat in late infancy. *Pediatr Obes*. 2019; 14(3). <https://doi.org/10.1111/ijpo.12476> PMID: 30362284
72. Sundholm JKM, Litwin L, Rönö K, Koivusalo SB, Eriksson JG, Sarkola T. Maternal obesity and gestational diabetes: Impact on arterial wall layer thickness and stiffness in early childhood—RADIEL study six-year follow-up. *Atherosclerosis*. 2019. <https://doi.org/10.1016/j.atherosclerosis.2019.01.037> PMID: 30819513
73. Scherrer U, Rimoldi SF, Rexhaj E, Stuber T, Duplain H, Garcin S, et al. Systemic and pulmonary vascular dysfunction in children conceived by assisted reproductive technologies. *Circulation*. 2012; 125(15):1890–1896. <https://doi.org/10.1161/CIRCULATIONAHA.111.071183> PMID: 22434595
74. de Arriba A, Domínguez M, Labarta JI, Puga B, Mayayo E, Longás AF. Metabolic syndrome and endothelial dysfunction in a population born small for gestational age relationship to growth and Gh therapy. *Pediatr Endocrinol Rev*. 2013; 10(3):297–307. PMID: 23724437
75. Maurice RL, Vaujois L, Dahdah N, Chibab N, Maurice A, Nuyt AM, et al. Carotid wall elastography to assess midterm vascular dysfunction secondary to intrauterine growth restriction: Feasibility and comparison with standardized intima-media thickness. *Ultrasound Med Biol*. 2014; 40(5):864–870. <https://doi.org/10.1016/j.ultrasmedbio.2013.11.013> PMID: 24495436
76. Xu GF, Zhang JY, Pan HT, Tian S, Liu ME, Yu TT, et al. Cardiovascular dysfunction in offspring of ovarian-hyperstimulated women and effects of estradiol and progesterone: A retrospective cohort study and proteomics analysis. *J Clin Endocrinol Metab*. 2014; 99(12):E2494–E2503. <https://doi.org/10.1210/jc.2014-2349> PMID: 25268391
77. Putra ST, Mansyur M, Sastroasmoro S. Effects of duration of breastfeeding during infancy on vascular dysfunction in adolescents. *Acta Med Indones*. 2015; 47(1):24–30. PMID: 25948764
78. Sodhi KS, Hondappanavar A, Saxena AK, Dutta S, Khandelwal N. Intima-media complex thickness: Preliminary workup of comparative evaluation of abdominal aorta and carotid artery of small-for-gestation-age term newborns and normal size term newborns. *Acta Cardiol*. 2015; 70(3):351–357. <https://doi.org/10.1080/ac.70.3.3080640> PMID: 26226709
79. Ciccone MM, Cortese F, Gesualdo M, DIM A, Tafuri S, Mancini G, et al. The role of very low birth weight and prematurity on cardiovascular disease risk and on kidney development in children: a pilot study. *Minerva Pediatr*. 2016. <https://doi.org/10.23736/S0026-4946.16.04520-5> PMID: 27355157
80. Faienza MF, Brunetti G, Delvecchio M, Zito A, de Palma FD, Cortese F, et al. Vascular function and myocardial performance indices in children born small for gestational age. *Circ J*. 2016; 80(4):958–963. <https://doi.org/10.1253/circj.CJ-15-1038> PMID: 26861187
81. Olander RFW, Sundholm JKM, Ojala TH, Andersson S, Sarkola T. Neonatal Arterial Morphology Is Related to Body Size in Abnormal Human Fetal Growth. *Circ Cardiovasc Imaging*. 2016; 9(9). <https://doi.org/10.1161/CIRCIMAGING.116.004657> PMID: 27601367
82. Dilli D, Ozkan E, Ozkan MB, Aydin B, Özyazici A, Fettah N, et al. Umbilical cord asymmetric dimethylarginine levels and ultrasound assessment of carotid arteries in neonates born small for gestational age. *J Matern Fetal Neonatal Med*. 2017; 30(4):492–496. <https://doi.org/10.1080/14767058.2016.1176136> PMID: 27072784
83. Wilde MAD, Eising JB, Gunning MN, Koster MPH, Evelein AMV, Dalmeijer GW, et al. Cardiovascular and Metabolic Health of 74 Children From Women Previously Diagnosed With Polycystic Ovary Syndrome in Comparison With a Population-Based Reference Cohort. *Reprod Sci*. 2018; 25(10):1492–1500. <https://doi.org/10.1177/1933719117749761> PMID: 29320957

84. Muñiz Fontán M, Oulego Erroz I, Revilla Orias D, Muñoz Lozón A, Rodríguez Núñez A, Lurbe IFE. Thoracic Aortic Intima-Media Thickness in Preschool Children Born Small for Gestational Age. *J Pediatr*. 2019. <https://doi.org/10.1016/j.jpeds.2018.12.037> PMID: 30732998
85. Crispi F, Figueras F, Cruz-Lemini M, Bartrons J, Bijmens B, Gratacos E. Cardiovascular programming in children born small for gestational age and relationship with prenatal signs of severity. *Am J Obstet Gynecol*. 2012; 207(2):121.e1–e9. <https://doi.org/10.1016/j.ajog.2012.05.011> PMID: 22717268
86. Valenzuela-Alcaraz B, Crispi F, Bijmens B, Cruz-Lemini M, Creus M, Sitges M, et al. Assisted reproductive technologies are associated with cardiovascular remodeling in utero that persists postnatally. *Circulation*. 2013; 128(13):1442–1450. <https://doi.org/10.1161/CIRCULATIONAHA.113.002428> PMID: 23985787
87. Meister TA, Rimoldi SF, Soria R, von Arx R, Messerli FH, Sartori C, et al. Association of Assisted Reproductive Technologies With Arterial Hypertension During Adolescence. *J Am Coll Cardiol*. 2018; 72(11):1267–1274. <https://doi.org/10.1016/j.jacc.2018.06.060> PMID: 30190005
88. Bianco-Miotto T, Craig JM, Gasser YP, van Dijk SJ, Ozanne SE. Epigenetics and DOHaD: from basics to birth and beyond. *J Dev Orig Health Dis*. 2017; 8(5):513–519. <https://doi.org/10.1017/s2040174417000733> PMID: 28889823
89. Allen NB, Krefman AE, Labarthe D, Greenland P, Juonala M, Kahonen M, et al. Cardiovascular Health Trajectories From Childhood Through Middle Age and Their Association With Subclinical Atherosclerosis. *JAMA Cardiol*. 2020. <https://doi.org/10.1001/jamacardio.2020.0140> PMID: 32159727
90. Belbasis L, Savvidou MD, Kanu C, Evangelou E, Tzoulaki I. Birth weight in relation to health and disease in later life: an umbrella review of systematic reviews and meta-analyses. *BMC Med*. 2016; 14(1):147. <https://doi.org/10.1186/s12916-016-0692-5> PMID: 27677312
91. Beune IM, Bloomfield FH, Ganzevoort W, Embleton ND, Rozance PJ, van Wassenaer-Leemhuis AG, et al. Consensus Based Definition of Growth Restriction in the Newborn. *J Pediatr*. 2018; 196:71–6.e1. <https://doi.org/10.1016/j.jpeds.2017.12.059> PMID: 29499988
92. Verburg BO, Jaddoe VW, Wladimiroff JW, Hofman A, Witteman JC, Steegers EA. Fetal hemodynamic adaptive changes related to intrauterine growth: the Generation R Study. *Circulation*. 2008; 117(5):649–659. <https://doi.org/10.1161/CIRCULATIONAHA.107.709717> PMID: 18212281
93. de Jong F, Monuteaux MC, van Elburg RM, Gillman MW, Belfort MB. Systematic review and meta-analysis of preterm birth and later systolic blood pressure. *Hypertension*. 2012; 59(2):226–234. <https://doi.org/10.1161/HYPERTENSIONAHA.111.181784> PMID: 22158643
94. Markopoulou P, Papanikolaou E, Analytis A, Zoumakis E, Siahianidou T. Preterm Birth as a Risk Factor for Metabolic Syndrome and Cardiovascular Disease in Adult Life: A Systematic Review and Meta-Analysis. *J Pediatr*. 2019; 210:69–80.e5. <https://doi.org/10.1016/j.jpeds.2019.02.041> PMID: 30992219
95. Juonala M, Cheung MM, Sabin MA, Burgner D, Skilton MR, Kahonen M, et al. Effect of birth weight on life-course blood pressure levels among children born premature: the Cardiovascular Risk in Young Finns Study. *J Hypertens*. 2015; 33(8):1542–1548. <https://doi.org/10.1097/HJH.0000000000000612> PMID: 26136063
96. Di Bernardo S, Mivelaz Y, Epure AM, Vial Y, Simeoni U, Bovet P, et al. Assessing the consequences of gestational diabetes mellitus on offspring's cardiovascular health: MySweetHeart Cohort study protocol. *BMJ Open*. 2017; 7(11):e016972. <https://doi.org/10.1136/bmjopen-2017-016972> PMID: 29138200
97. van den Oord SC, Sijbrands EJ, ten Kate GL, van Klaveren D, van Domburg RT, van der Steen AF, et al. Carotid intima-media thickness for cardiovascular risk assessment: systematic review and meta-analysis. *Atherosclerosis*. 2013; 228(1):1–11. <https://doi.org/10.1016/j.atherosclerosis.2013.01.025> PMID: 23395523
98. Sundholm JK, Olander RF, Ojala TH, Andersson S, Sarkola T. Feasibility and precision of transcutaneous very-high resolution ultrasound for quantification of arterial structures in human neonates—comparison with conventional high resolution vascular ultrasound imaging. *Atherosclerosis*. 2015; 239(2):523–527. <https://doi.org/10.1016/j.atherosclerosis.2015.02.016> PMID: 25721703
99. Sundholm JKM, Paetau A, Albäck A, Pettersson T, Sarkola T. Non-Invasive Vascular Very-High Resolution Ultrasound to Quantify Artery Intima Layer Thickness: Validation of the Four-Line Pattern. *Ultrasound Med Biol*. 2019; 45(8):2010–2018. <https://doi.org/10.1016/j.ultrasmedbio.2019.04.017> PMID: 31101444
100. Mayanskaya SD, Grebenkina IA. Structural and Functional Remodeling of Vascular Wall in Persons With Family History of Hypertension. *Kardiologiia*. 2015; 55(2):27–31. PMID: 28294807
101. Pac-Kozuchowska E, Zatorska-Karpuś M, Wieczorek P. Comparison of lipid and lipoprotein(a) metabolism in blood serum and the thickness of common carotid artery (intima-media thickness complex—

- IMT), in children from families at risk for cardiovascular disease and in families without risk. *Med Wieku Rozwoj*. 2005; 9(2):171–178. PMID: [16085958](#)
102. Arnaiz P, Pino F, Marín A, Barja S, Aglony M, Cassis B, et al. Association between a cardiovascular risk score and early markers of atherosclerotic disease in Chilean children. *Rev Med Chil*. 2010; 138(10):1226–1231. <https://doi.org/S0034-98872010001100003> PMID: [21279267](#)
 103. Lee AC, Kozuki N, Cousens S, Stevens GA, Blencowe H, Silveira MF, et al. Estimates of burden and consequences of infants born small for gestational age in low and middle income countries with INTERGROWTH-21(st) standard: analysis of CHERG datasets. *BMJ*. 2017; 358:j3677. <https://doi.org/10.1136/bmj.j3677> PMID: [28819030](#)
 104. Wollmann HA. Intrauterine growth restriction: definition and etiology. *Horm Res*. 1998; 49(Suppl 2):1–6. <https://doi.org/10.1159/000053079> PMID: [9716819](#)
 105. Chamberlain C, O'Mara-Eves A, Porter J, Coleman T, Perlen SM, Thomas J, et al. Psychosocial interventions for supporting women to stop smoking in pregnancy. *Cochrane Database Syst Rev*. 2017; 2. <https://doi.org/10.1002/14651858.CD001055.pub5> PMID: [28196405](#)
 106. Barker M, Dombrowski SU, Colbourn T, Fall CHD, Kriznik NM, Lawrence WT, et al. Intervention strategies to improve nutrition and health behaviours before conception. *Lancet*. 2018; 391(10132):1853–1864. [https://doi.org/10.1016/S0140-6736\(18\)30313-1](https://doi.org/10.1016/S0140-6736(18)30313-1) PMID: [29673875](#)
 107. Hanson M, Gluckman P. Commentary: Developing the future: life course epidemiology, DOHaD and evolutionary medicine. *Int J Epidemiol*. 2016; 45(4):993–996. <https://doi.org/10.1093/ije/dyw105> PMID: [27880687](#)
 108. Falkner B, Lurbe E. Primordial Prevention of High Blood Pressure in Childhood: An Opportunity Not to be Missed. *Hypertension*. 2020; 75(5):1142–1150. <https://doi.org/10.1161/HYPERTENSIONAHA.119.14059> PMID: [32223379](#)